

BY 3 DAYS OF AGE:

SUPPLEMENTATION with IRON:

Ferroforte (comes in 100mL bottle) – **1mL DOSE**

- **Piglets:**
 - ALL piglets to get one dose no later than 3 days of age



AT WEANING:

VACCINATION against WASTING DISEASES:

Circo Virus

Circoflex (comes in 50 dose bottles) – **1mL DOSE**

21 day withdrawal

- **Piglets:**
 - One dose ONCE at 3 weeks of age



AT WEANING AND 3-4 WEEKS LATER:

VACCINATION against SOIL-BORNE disease:

Erysipelas

ER Bac Plus (comes in 50 dose bottles) – **2mL DOSE**

21 day withdrawal

- **Piglets:**
 - 2ml dose at >4 weeks of age
 - Booster 2ml dose 3-4 weeks after first dose



VACCINATION against SOIL-BORNE disease:

Clostridial Disease

Ultrachoice 8 (comes in 50 dose bottles) – **2mL DOSE**

21 day withdrawal

- **Piglets:**
 - 2ml dose at weaning
 - Booster 2ml dose 3-4 weeks after first dose



For more information on best practices for raising pigs outdoors, refer to the *Canadian Small-Scale Pig Farming Manual*: www.casv-acvp.com/small-scale-pig-farming.html

AROUND 6-10 WEEKS OF AGE AND 4-5 WEEKS BEFORE SLAUGHTER:

DEWORMING:

Safeguard (does not treat mites or threadworms)

0 day withdrawal

OR

Ivomec (does not treat trichuris suis, but does treat mange/mites)

28 day withdrawal (injectable)

14 day withdrawal (top dress)

- **Finishing Pigs:**

- Treat between 6 and 10 weeks old
- Treat 4-5 weeks prior to slaughter (about 21 to 25 weeks)



If you find milk spot livers after one turn of this deworming protocol, add in a 3rd deworming dose in the finisher pigs (about week 15/16)!!!

******Note: you can rotate dewormers******

